

Vaccination Policy

At Bright Oaks Pediatric Center, we strive to provide the best care that is based on the most up-to-date and accurate literature for our patients and families. In order to do this, we require all of our patients to receive immunizations on schedule, without spacing, delaying, alternate schedules, or otherwise.

Our vaccine schedule is noted on the following pages.

While we respect one's choice to not vaccinate, we feel we would be doing a disservice to our patients and community, and feel uncomfortable as a practice caring for children who do not receive these recommended immunizations at all or whose families choose to go off schedule. The risk for errors and incomplete immunization increases any time there is deviation from the routine vaccination schedule.

Vaccines are a simple and easy way to prevent deadly illnesses and diseases. We ask that should you have questions or concerns regarding immunizations, ask your child's healthcare provider. Unfortunately, there is a lot of misinformation on the internet, as anyone can publish or post an article. When researching, ensure you are looking at reputable and quality resources.

If a vaccine or vaccines have already been drawn up at your child's scheduled appointment and are not administered at the visit, you will be responsible to pay for the full cost of the vaccine(s) as we are unable to bill insurance companies for vaccines not administered to a patient.

We look forward to caring for your child at Bright Oaks!



RECOMMENDED IMMUNIZATIONS AND PHYSICAL EXAM SCHEDULE

| AGE | IMMUNIZATIONS | PHYSICAL EXAM PROCEDURES |
|------------------------|--|--|
| Birth | Hepatitis B #1 (done in hospital) | NEWBORN SCREENING #1, HEARING SCREEN (in hospital) |
| Newborn Appointment | Hepatitis B #1 (if not done in hospital) | *Not considered a wellness visit by insurance companies |
| 2 Weeks | | NEWBORN SCREENING #2 (heel stick), EPDS (Edinburgh Postnatal Depression Scale) |
| 4 Weeks | | EPDS |
| 2 Months | Vaxelis, Prevnar, Rotavirus(oral vaccine) | EPDS |
| 4 Months | Vaxelis, Prevnar, Rotavirus(oral vaccine) | EPDS |
| 6 Months | Vaxelis, Prevnar, Rotavirus(oral vaccine) *Influenza if in season (optional) | Lead Risk Assessment, EPDS |
| 9 Months | | Lead Risk Assessment |
| 12 Months | MMR, Varicella, Hepatitis A #1 | Lead & Hemoglobin Test (fingerstick), TB Risk Assessment, Lead Risk Assessment, Photoscreen (vision) |
| 15 Months | Pentacel & Prevnar | TB Risk Assessment, Lead Risk Assessment |
| 18 Months | Hepatitis A #2 | Developmental Screen TB Risk Assessment, Lead Risk Assessment |
| 2 Years | | Developmental Screen, Lead Test (fingerstick), TB Risk Assessment, Lead Risk Assessment, Cholesterol Risk Assessment, Photoscreen (vision) |
| 2 ½ Years | | |



2111 Laurel Bush Rd. Ste H Bel Air, MD 21015 (p) 410-569-3300 (f) 410-569-8199 www.BrightOaksPediatrics.com

| | Padiatric Center | <u>www.BrightOaksPediatrics.com</u> |
|-------------|---------------------------------------|--|
| | | Photoscreen (vision), TB Risk Assessment, |
| 3 Years | | Lead Risk Assessment, |
| 0.000 | | Cholesterol Risk Assessment |
| | | Choloster of Talox 7 lococomons |
| | ProQuad & Quadracel | Vision & Hearing Screen, TB Risk Assessment, |
| 4 Years | | Lead Risk Assessment, |
| 7 10013 | | Cholesterol Risk Assessment |
| | | Cholosteror Mak / Addedmank |
| | | Vision & Hearing Screen, Cholesterol Risk |
| 5 Years | | Assessment, TB Risk Assessment, Lead Risk |
| Jicais | | Assessment Assessment |
| | | Assessment |
| | | TB Risk Assessment, Lead Risk Assessment, |
| 6 Years | | Cholesterol Risk Assessment, |
| o rears | | · |
| | | Vision and Hearing |
| | | TB Risk Assessment, |
| 7 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| | | |
| | | TB Risk Assessment, |
| 8 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| 0 Tears | | The state of the s |
| | | TB Risk Assessment, |
| 9 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| 9 Teals | | The state of the s |
| | | TB Risk Assessment, |
| 10 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| 10 lears | | The state of the s |
| | Tdap & Meningitis ACWY #1 | TB Risk Assessment, |
| 11 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| ii feais | *HPV #1(optional) | Cholesteror Risk Assessment, Vision and Hearing |
| | *110\/ #2/#:1\ | TD Diek Assessment |
| | *HPV #2(optional) | TB Risk Assessment, |
| 12-14 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| | | TD Diale Assessment |
| | | TB Risk Assessment, |
| 15 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| | Maningitia ACVAN/#0 9 Maningitia D #4 | TD Diek Assessment |
| 40.14 | Meningitis ACWY #2 & Meningitis B #1 | TB Risk Assessment, |
| 16 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| | | |
| | Meningitis B #2 | TB Risk Assessment, |
| 17-18 Years | | Cholesterol Risk Assessment, Vision and Hearing |
| | | |

The Influenza Vaccine is recommended each year for children 6 months and older.



2111 Laurel Bush Rd. Ste H Bel Air, MD 21015 (p) 410-569-3300 (f) 410-569-8199 www.BrightOaksPediatrics.com

Vaccine Key:

- **Hepatitis B** Protects against infection from the Hepatitis B virus (HBV) and prevents potential complications like chronic liver disease and liver cancer.
- **Hepatitis A** Protects against infection from the Hepatitis A virus (a serious liver infection).
- · **Vaxelis** Protects against Diphtheria, Tetanus, Pertussis (whooping cough), Polio, Haemophilus Influenzae type b, Hepatitis b.
- Rotavirus Protects against Rotavirus, a contagious gastrointestinal (GI) infection.
- **Prevnar** Protects against bacteria that cause invasive Pneumococcal disease (IPD).
- **Pentacel –** Protects against Diphtheria, Tetanus, Pertussis (whooping cough), Polio, Haemophilus Influenzae type b.
- MMR Protects against Measles, Mumps and Rubella.
- Varicella Protects against Chickenpox.
- ProQuad Protects against Measles, Mumps, Rubella, and Varicella.
- Quadracel Protects against Diphtheria, Tetanus, Pertussis (whooping cough), Polio.
- **Tdap** Protects against Tetanus, Diphtheria, and Pertussis (whooping cough).
- · **Meningitis ACWY** Protects against four types of Meningococcal bacteria: A, C, W, and Y, which can cause life-threatening illnesses like meningitis (inflammation of the membranes covering the brain and spinal cord) and sepsis.
- **Meningitis B** Protects against serious and potentially fatal infections caused by serogroup B Meningococcal bacteria. This includes meningitis (inflammation of the membranes covering the brain and spinal cord) and blood infections.
- *HPV Protects against the types of Human Papillomaviruses that cause most cervical, vaginal, vulvar, penile, anal, and oropharyngeal (head and neck) cancers, as well as genital warts. (Two doses if started before age 15 and three doses if started at age 15 or later.) *This vaccine is optional.
- *Influenza Protects against different strains of Influenza A & B. *This vaccine is optional.